







HEAD LICE-PRINCIPLES TO GUIDE SCHOOL POLICIES

USEFUL INFORMATION

- * One of the worst “problems” of head lice is adult attitudes; having head lice is not a serious medical condition.
- * “Irrational” reactions to head lice, e.g., fumigating classrooms, school buses, etc., are expensive and unnecessary.
- * Much “information” about head lice is based on old, unproven information generated more than 80 years ago, some of it propagated by the companies which sell lice shampoos and sprays.
- * It is possible but difficult to obtain head lice from casual head to head contact.
- * Transmission via clothing, hats, furniture, carpets, is also possible.
- * Lice are fragile, but can be passed on by hats and combs in some cases.
- * There is no significant relationship between hair length or personal cleanliness and transmission.
- * It is unlikely that a nit on a stray hair shaft will hatch because the optimal conditions for hatching exist only on the human head.
- * Stray lice that fall off a head are injured or dying and incapable of causing a new infestation.
- * In time, inbreeding of lice on a person’s head causes them to die spontaneously; that’s why kids do not become covered with them. It is a self-limiting condition.
- * It is possible to tell whether treatment has been successful by the appearance of the eggs.
- * **CONSIDERING THE AVERAGE CASE OF HEAD LICE IS 3-4 MONTHS OLD BEFORE IT IS DETECTABLE, A STRICT NO NIT POLICY IS NOT NECESSARY AND ONLY DEPRIVES CHILDREN OF EDUCATIONAL TIME.**
- * Although schools, day care centers, etc. are often blamed for head lice outbreaks, it is the family unit that maintains cases leading to outbreaks in schools.

Health Office
182 Thurbers Avenue
Providence, RI 02905
tefTT3 1 77 0.24 0 0 0.24 519.0263